

# The Optical Afterglow of GRB 971214: R and J Photometry

A. Diercks<sup>1</sup>, E. W. Deutsch<sup>1</sup>, F. J. Castander<sup>2</sup>, C. Corson<sup>3</sup>, G. Gilmore<sup>4</sup>, D. Q. Lamb<sup>2</sup>,  
N. Tanvir<sup>4</sup>, E. L. Turner<sup>5</sup>, and R. Wyse<sup>6</sup>

Accepted for publication in The Astrophysical Journal Letters

Volume 503, L105

*received 1998 March 23; accepted 1998 June 24*

## ABSTRACT

We present an *R*-band and *J*-band photometry of an optical transient which is likely to be associated with the gamma-ray burst event GRB 971214. Our first measurement took place 13 hours after the gamma-ray event. The brightness decayed with a power-law exponent  $\alpha = -1.20 \pm 0.02$ , which is similar to those of GRB 970228 and GRB 970508 which had exponents of  $\alpha = -1.10 \pm 0.04$  and  $\alpha = -1.141 \pm 0.014$  respectively. The transient decayed monotonically during the first four days following the gamma-ray event in contrast with the optical transient associated with GRB 970508 which increased in brightness, peaking two days after the burst, before settling to a power-law decay.

*Subject headings:* gamma rays: bursts

---

<sup>1</sup>Department of Astronomy, Box 351580, University of Washington, Seattle, WA 98195; diercks@astro.washington.edu; deutsch@astro.washington.edu

<sup>2</sup>Department of Astronomy and Astrophysics, University of Chicago, 5640 South Ellis Avenue, Chicago, IL 60637; fjc@oddjob.uchicago.edu; lamb@pion.uchicago.edu

<sup>3</sup>Apache Point Observatory, 2001 Apache Point Rd., Sunspot NM 88349; corson@galileo.apo.nmsu.edu

<sup>4</sup>Institute of Astronomy, Madingley Road, Cambridge CB3 0HA, UK; gil@ast.cam.ac.uk

<sup>5</sup>Princeton University Observatory, Peyton Hall, Princeton, NJ 08544; elt@astro.princeton.edu

<sup>6</sup>Department of Physics and Astronomy, The Johns Hopkins University, Baltimore, MD 21218; wyse@wyser.pha.jhu.edu

## 1. Introduction

The launch of the BeppoSAX satellite (Boella et al. 1997) in 1996 has led to a recent breakthrough in the study of gamma-ray bursts by detecting fading X-ray counterparts and localizing them to a few arc-minutes on the sky. This has allowed subsequent identification of optical counterparts in three cases, GRB 970228 (Groot et al. 1997), GRB 970508 (Bond 1997), and GRB 971214 (Halpern et al. 1997).

GRB 971214 triggered the BeppoSAX Gamma Ray Burst Monitor on December 14.97272 UT with a peak flux of  $650 \text{ counts s}^{-1}$ . In addition, the burst was localized to an error radius of  $3'9$  (99% confidence) with the Wide Field Camera (Heise et al. 1997). The gamma-ray event was also detected by BATSE which measured a total fluence above 20 keV of  $1.09 \pm 0.07 \times 10^{-5} \text{ erg cm}^{-2}$  and by one RXTE-ASM camera yielding a peak intensity of  $470 \pm 140 \text{ mCrab}$  (Kippen et al. 1997). Shortly thereafter, a fading optical source was detected within the BeppoSAX error circle at  $\alpha(J2000) = 11^{\text{h}}56^{\text{m}}26^{\text{s}}.4$ ,  $\delta(J2000) = 65^{\circ}12'00''.5$  with I-band magnitudes of  $21.2 \pm 0.3$  on Dec 15.47 UT and  $\sim 22.6$  on Dec 16.47 UT (Halpern et al. 1997). Further observations by BeppoSAX detected a previously unknown fading X-ray source, later designated 1SAX J1156.4+6513, within the initial error circle at  $\alpha(J2000) = 11^{\text{h}}56^{\text{m}}25^{\text{s}}$ ,  $\delta(J2000) = 65^{\circ}13'11''$  with an error radius of about  $1'$  (Antonelli et al. 1997). Since this second X-ray detection is consistent with the position of the fading optical source identified by Halpern et al. (1997) it is quite likely that these objects are the X-ray and optical afterglow of GRB 971214.

We report here *R*-band and *J*-band observations of this optical transient (OT) made with the Apache Point Observatory (APO) 3.5 m telescope. This work extends the preliminary photometry reported in Diercks et al. (1997), Castander et al. (1997), and Tanvir et al. (1997).

## 2. *R*-band Observations

The field of GRB 971214 was imaged on the four consecutive nights following the burst, with the first observations taking place 13 hours after the BeppoSAX detection. The optical transient was positively detected each night. All observations were taken with a thinned SITe 2048 CCD through an *R*-band filter at  $2 \times 2$  binning yielding a plate scale of  $0''.28 \text{ pixel}^{-1}$ . The images were overscan subtracted and flat-fielded with twilight flats in the usual manner using IRAF (Tody 1993). The field of GRB 971214 is plagued by the presence of a bright star HD103690 ( $V = 6.7$ ) which contributed a substantial amount of stray light to the images. In order to optimally combine the images, we removed this stray light by median filtering each frame with a  $14''$  rectangular box and then smoothing the resulting image with a  $18''$  FWHM Gaussian filter. The spatially varying part of each smoothed image was then subtracted from its corresponding flat-fielded image to remove the excess stray light. As the variations were typically less than 10%, the effect on the noise computations due to slightly lower sky values in certain regions of the images was quite small.

All the images for each night were aligned and combined, weighting each frame by the photon noise within a seeing disk. The epoch quoted for each measurement is average of the time of each observation weighted in the same manner. The four final images are shown in Figure 1.

The OT discovered by Halpern is clearly visible in our December 15.51 data and has sufficient signal to fit an accurate centroid using DoPhot (Schechter, Mateo, & Saha 1993). We measure a position of  $\alpha(J2000) = 11^{\text{h}}56^{\text{m}}26^{\text{s}}.6$ ,  $\delta(J2000) = 65^{\circ}12'00''.9$  for the OT with an uncertainty of  $0''.5$  using five nearby stars in the USNO-A1.0 catalog (Monet 1996). All subsequent photometry on both the OT and comparison stars was done using DoPhot with these positions fixed as it resulted in greater precision especially in the later data where the OT is quite faint.

In order to test the robustness of our detection in the December 18.50 UT data where the OT is close to the detection limit, we performed fixed position PSF photometry on 50 arbitrary regions within the image. In only one case was there sufficient signal to fit a PSF, and the returned magnitude was at the detection limit of the image. We are therefore confident in our detection of the OT in the December 18.50 data.

We observed the field PG0918+029 (Landolt 1992) and the field of GRB971214 on 1998 February 1 under photometric conditions with the same camera and filter arrangement used to monitor the OT. Observations were made over a range of airmass from 1.18 to 2.32 and we derived an average R-band extinction of  $0.088 \pm 0.031 \text{ mag airmass}^{-1}$ .

Using this coefficient we obtained calibrated magnitudes for five stars in the field of GRB971214 which are listed in Table 1 and shown in Figure 1. Star 3 was saturated in some of the monitoring data and was not used for calibration purposes. Using the remaining stars, we normalized the monitoring data to obtain the absolute photometry of the OT on each night. The lightcurve for the four days following the burst is shown in Figure 2 along with the lightcurves of the OTs associated with GRB 970228 (Galama et al. 1997) and GRB 970508 (Galama et al. 1998) for comparison.

### 3. J-band Observations

We also imaged the GRB971214 field with the Grism Spectrometer and Imager II (GRIM II) in the J band on the first two nights after the burst. The GRIM II instrument has a NICMOS array, which at f/5 provides a pixel scale of  $0.47''$  and a field of view of  $2' \times 2'$ . We obtained dithered 10-second exposures, totaling 730 and 1150 seconds on the first and second nights, respectively. Images were dark-current and bad-pixel corrected, flat-fielded and combined, with high sigma threshold clipping, into a single image for each night. The mean time of the resulting exposures were December 15.44 and 16.45 UT, respectively.

We detect the OT at a magnitude of  $J = 20.47_{-0.19}^{+0.21}$  on December 15.44 and  $J = 21.46_{-0.26}^{+0.34}$  on December 16.45, where the calibration is performed using observations of the UKIRT faint standard star FS12, carried out at several airmasses. Table 2 summarizes both the R-band and J-band observations.

### 4. Discussion

The well-observed light curve of the GRB970508 OT, the brightest observed thus far, shows a dramatic rise, peaking nearly two days after the initial burst, before beginning a power-law

decay. GRB 970228 was not observed nearly as often through a consistent filter, but there is also evidence (depending on spectral assumptions) (Guarnieri et al. 1997) that the transient increased in brightness until  $\sim 20$  hr after the burst, after which it also began fading with a power-law slope essentially identical to that of GRB 970508. Despite the same decay slope, the GRB 970228 OT was  $\sim 1.5$  mag fainter than the GRB 970508 OT. A detailed analysis of the difference between these two light curves is presented in Pedersen et al. (1998).

A power-law of the form  $F = F_0 t^\alpha$  was fit to the four  $R$ -band data points yielding  $\alpha = -1.20 \pm 0.02$ . This rate of decline is similar to the two previously identified bursts although there is no evidence of the OT brightening over the course of the observations.

The observations from all four nights are combined into one deep image totaling 3.25 hrs of integration with a limiting  $R$ -band magnitude  $\sim 25.4$  (Figure 3). The two brightest objects within  $20''$  of the OT are an extended source (A)  $4''.6$  southwest of the transient with  $R = 22.7 \pm 0.1$ , and an unresolved source (B)  $4''.9$  north of the transient with  $R = 23.43 \pm 0.08$ . The resolution of these images is insufficient to identify any structure in the extended object.<sup>7</sup>

## 5. Acknowledgments

We would like to thank Jan van Paradjis and the BeppoSAX team for their critical help in making our early observations of GRB 971214 possible. We would also like to thank Bruce Margon, Christopher Stubbs, and Eli Waxman for useful and informative discussions. These observations were made on very short notice and involved last-minute technical changes which would not have been possible without the enthusiastic cooperation of the APO site staff. We thank Daniel Reichart for pointing out an error in the UT times of the APO  $R$ -band observations in a pre-print of this paper, which when corrected, yield a slower rate of decline. AD acknowledges the generous support of the David and Lucille Packard Foundation.

---

<sup>7</sup>To obtain the images discussed in this work, contact A. Diercks or E. Deutsch

Table 1. Calibration stars used in the field of GRB 971214. The uncertainties reflect photon statistics as well as uncertainty in the determination of the zero-point.

| Star | $\alpha$ (J2000) | $\delta$ (J2000) | $R$              |
|------|------------------|------------------|------------------|
| 1    | 11 56 25.8       | 65 11 36.3       | $19.96 \pm 0.03$ |
| 2    | 11 56 34.2       | 65 11 44.4       | $20.66 \pm 0.03$ |
| 3    | 11 56 29.8       | 65 12 16.3       | $16.38 \pm 0.03$ |
| 4    | 11 56 20.1       | 65 12 34.6       | $20.38 \pm 0.03$ |
| 5    | 11 56 13.4       | 65 11 15.6       | $20.50 \pm 0.03$ |

Table 2. Summary of APO observations of the GRB 971214 optical afterglow.

| Date (UT) | Days Since Burst | Exposures | Seeing (arcsec) | $2\sigma$ Detection Limit | Magnitude $\pm 1\sigma$     |
|-----------|------------------|-----------|-----------------|---------------------------|-----------------------------|
| Dec 15.44 | 0.47             | 730s      | 1.3             | 21.8                      | $J = 20.47^{+0.21}_{-0.19}$ |
| Dec 15.51 | 0.54             | 3300s     | 1.4             | 24.6                      | $R = 22.06 \pm 0.06$        |
| Dec 16.45 | 1.48             | 1150s     | 1.7             | 21.6                      | $J = 21.46^{+0.34}_{-0.26}$ |
| Dec 16.53 | 1.56             | 1800s     | 1.2             | 24.4                      | $R = 23.36 \pm 0.13$        |
| Dec 17.50 | 2.53             | 3000s     | 1.2             | 24.8                      | $R = 24.08 \pm 0.16$        |
| Dec 18.50 | 3.53             | 3600s     | 1.1             | 24.8                      | $R = 24.61 \pm 0.22$        |

## REFERENCES

- Antonelli, L. A., Butler, R. C., Piro, L., Celidonio, G., Coletta, A., Tesseri, DeLibero, C. 1997, IAU Circ. 6792
- Band, D. L. 1997, in The Eighth Marcel Grossmann Meeting on General Relativity (Singapore: World Scientific), in press, (astro-ph/9712193)
- Boella, G., Butler, R. C., Perola, G. C., Piro, L., Scarsi, L., & Bleeker, J. 1997, A&AS, 122, 299
- Bond, H. E. 1997, IAU Circ. 6654
- Castander, F. J., Wyse, R., Gilmore, G., Corson, C., Diercks, A., Deutsch, E. W., Lamb, D. Q., & Turner, E. 1997, IAU Circ. 6791
- Castro-Tirado, J. A. 1998, in ESA Conference Proceedings SP-41, Ultraviolet Astrophysics: beyond the IUE Final Archive, ed. R. Gonzalez-Riestra, W. Wamsteker, & R. Harris (Noordwijk: ESA Publications Division), 659
- Diercks, A., Deutsch, E. W., Wyse, R., Gilmore, G., Corson, C., Castander, F. J. & Turner, E. 1997, IAU Circ. 6791
- Fishman, G. J. & Meegan, C. A. 1995, ARA&A, 33, 415
- Groot, P. J. et al. 1997, IAU Circ. 6584
- Galama, T. J. et al. 1998, in Gamma-Ray Bursts, 4-th Huntsville Symposium, ed. C. Meegan, R. Preece, & T. Koshut (New York: Springer-Verlag), in press, (astro-ph/9712322)
- Galama, T. J. et al. 1998, ApJL, in press, (astro-ph/9802160)
- Guarnieri, A. et al. 1997, A&A, 328, L13
- Halpern, J., Thorstensen, J., Helfand, D., & Costa, E. 1997, IAU Circ. 6788
- Heise, J. et al. 1997, IAU Circ. 6787
- Kippen, R. M., Woods, P., Connaughton, V., Smith, D. A., Levine, A. M., Remillard, R., Hurley, K. 1997, IAU Circ. 6789
- Landolt, A. U. 1992, AJ, 104, 340
- Monet, D. et al. 1996, USNO-A1.0 (Washington DC: US Naval Observatory)
- Pedersen, H. et al. 1998, ApJ 496, in press, (astro-ph/9710322)
- Schechter, P., Mateo, M., & Saha, A. 1993, PASP, 105, 1342
- Tanvir, N., Wyse, R., Gilmore, G., Corson, C. 1997, IAU Circ. 6796.
- Tody, D. 1993, in ASP Conf. Ser. 52, Astronomical Data Analysis Software and Systems II, ed. R.J. Hanish, R.J.V. Brissenden, & J. Barnes (San Francisco:ASP),173

The resolution of this figure is reduced from the original as requested by this archive.  
For a preprint with full resolution figures, see <http://www.astro.washington.edu/deutsch/pubs/>




Fig. 1.—  $100'' \times 100''$  R-band images of the field of GRB 971214 on the four nights observed from APO. The clearly fading optical transient is indicated with an arrow, and several of the comparison stars are numbered.




Fig. 2.—  $R$ -band lightcurve of the optical transient associated with GRB 971214 based on our observations at APO. The  $R$ -band lightcurve data points and power-law decay fit lines of the afterglows of GRB 970228 (*squares*) (Galama et al. 1997) and GRB 970508 (*triangles*) (Galama et al. 1998) are also shown for comparison. A thin dashed line is drawn between the data points measured during the rising phase of the GRB 970508 lightcurve.

The resolution of this figure is reduced from the original as requested by this archive.  
For a preprint with full resolution figures, see <http://www.astro.washington.edu/deutsch/pubs/>




Fig. 3.—  $150'' \times 120''$  R-band combined deep image of the GRB 971214 field. The optical transient is marked with an arrow. Photometric reference stars 1 – 5 and selected other objects discussed in the text are labeled.